

## **AIMVT Approved Abbreviation List**

The following are accepted abbreviations that do not require expansion in your application (both logs and reports):

<ul> <li>Ab</li> </ul>	antibody
------------------------	----------

ACT activated clotting timeAGID agar gel immunodiffusion

BID twice a day
BP blood pressure
bpm beats per minute
BUN blood urea nitrogen

BW body weight

C1, C2... cervical vertebraeC/M castrated male

CBC complete blood count

• cc cubic centimeter

cm centimeter

CNS central nervous system

CO2 carbon dioxide

CPD citrate-phosphate-dextrose

CPD-A citrate-phosphate-dextrose-adenine

CPK creatinine phosphokinaseCPR cardiopulmonary resuscitation

CSF cerebrospinal fluid

d daydL deciliter

DNA deoxyribonucleic acid

ECG/EKG electrocardiogram or electrocardiographic

EDTA ethylenediaminetetraacetic acid

ELISA enzyme-linked immunosorbent assay

°F degree Fahrenheit
EMG electromyography
F/S female/spayed
FeLV feline leukemia virus

FIP feline infectious peritonitisFIV feline immunodeficiency virus

g gram(s)
h hour
Hct hematocrit
Hgb hemoglobin
HPF high power field



hr hour(s)ID intradermal

IFA indirect fluorescent antibody

IM intramuscularIN intranasalin inch

IP intraperitonealIV intravenouskg kilogram

L1, L2... lumbar vertebrae

L literlb pound

LPF low power field

m meter

M/N male/neutered

MCH mean corpuscular hemoglobin

MCHC mean corpuscular hemoglobin concentration

MCV mean corpuscular volume

min minutemg milligramsmL milliliter

MRI magnetic resonance imaging
 NPO nothing by mouth (nil per os)

NSAID non-steroidal anti-inflammatory drug

• O2 oxygen

OD right eye (oculus dexter)OS left eye (oculus sinister)

OU both eyesoz ounce

PCV packed cell volumePE physical exam

• pH measure of the acidity of a solution

PMI point of maximum intensityPRN as required / as needed

PO per os
q every
QD once a day
QID four times a day
QOD every other day
RBC red blood cell

rDVM referring Doctor of Veterinary Medicine



RER resting energy requirement

RNA ribonucleic acid

• Rx take, receive – used to indicate a prescription or treatment

• s second

SC / SQ subcutaneous

SGOT serum glutamic-oxaloacetic transaminase
 SGPT serum glutamate-pyruvate transaminase

SID once a dayT4 thyroxine

T3 triiodothyronineTID three times a day

• TSH thyroid stimulating hormone

WBC white blood cell

wk weekwt weightyr year